



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$\text{Area } DEG = \pi r^2 (1 - \sin \theta \operatorname{cosec} \phi)^2.$$

$$u = x^2 (\phi \operatorname{cosec}^2 \phi - \cot \phi) = r^2 \sin^2 \theta (\phi \operatorname{cosec}^2 \phi - \cot \phi).$$

An element of surface at Q is $4x dx d\rho = 4r^2 \sin \theta \cos \theta d\theta d\rho$; at R it is $du = 2r^2 \sin^2 \theta \operatorname{cosec}^2 \phi (1 - \phi \cot \phi) d\phi$.

The limits of θ are 0 and $\frac{1}{2}\pi$; of ϕ , θ and $\pi - \theta$; of ρ , 0 and 2π .

Hence, doubling as R may be on either side of PQ , we get for the required chance,

$$\begin{aligned} p &= \frac{16\pi r^6}{\pi^3 r^6} \int_0^{\frac{1}{2}\pi} \int_{\theta}^{\pi-\theta} \int_0^{2\pi} \sin^3 \theta \cos \theta \operatorname{cosec}^2 \phi (1 - \phi \cot \phi) (1 - \sin \theta \operatorname{cosec} \phi)^2 d\theta d\phi d\rho \\ &= \frac{32}{\pi} \int_0^{\frac{1}{2}\pi} \int_{\theta}^{\pi-\theta} \sin^3 \theta \cos \theta \operatorname{cosec}^2 \phi (1 - \phi \cot \phi) (1 - \sin \theta \operatorname{cosec} \phi)^2 d\theta d\phi \\ &= \frac{2}{3\pi} \int_0^{\frac{1}{2}\pi} [2(\pi - 2\theta) \sin 2\theta + 4 - 4\cos 4\theta - 3\sin^2 2\theta \cos 2\theta \\ &\quad + 64\sin^4 \theta \cos \theta \log \tan \frac{1}{2}\theta] d\theta = \frac{2}{5}. \end{aligned}$$

Also solved by F. P. Matz.

Miscellaneous 146 was also solved by Jeannette Brooks.

PROBLEMS FOR SOLUTION.

ALGEBRA.

223. Proposed by THEODORE L. DELAND, Office of the Secretary of the Treasury, Washington, D.C.

An officer in the Treasury Department assigned three clerks to count a lot of silver dollars and when finished noted that there was an apparent difference in their efficiency; and, to determine the fact, gave to each a similar lot of the same amount to count, the only record made at the time being that A to count his lot alone, took three weeks longer, B took two weeks longer, and C took one week longer than it took for all working together to count the first lot. The best counter, on the record made, was given an efficiency mark of 93 on the scale of 100. What efficiency mark should, on the record, be given to each of the other two counters?

224. Proposed by G. W. GREENWOOD, M. A. (Oxon). Lebanon, Ill.

Show that, if none of the quantities x , y , z is zero, the result of eliminating them from

$$(x+y)(x+z) = bcyz \dots\dots (1),$$

$$(y+z)(y+x) = caxx \dots\dots (2),$$

$$(z+x)(z+y) = abxy \dots\dots (3),$$

$$\text{is } \begin{vmatrix} \pm a, & 1, & 1 \\ 1, & \pm b, & 1 \\ 1, & 1, & \pm c \end{vmatrix} = 0.$$

[Oxford, 1896.]

225. Proposed by H. M. ARMSTRONG, Cooch's Bridge, Delaware.

If $a = ax + cy + bz \dots\dots (1)$, $\beta = cx + by + az \dots\dots (2)$, $\gamma = bx + ay + cz \dots\dots (3)$, show that $a^3 + \beta^3 + \gamma^3 - 3a\beta\gamma = (a^3 + b^3 + c^3 - 3abc)(x^3 + y^3 + z^3 - 3xyz)$.

GEOMETRY.

248. Proposed by CHRISTIAN HORNING, Heidelberg University, Tiffin, Ohio.

Given AB, BC in a straight line, to produce it to D so that $AD \cdot CD = BD^2$.

249. Proposed by W. W. BEMAN, The University of Michigan.

Given the distances of a point in the plane of a square from three of its vertices, to find the side of the square.

250. Proposed by W. W. BEMAN, The University of Michigan.

Given the distances of a point in the plane of an equilateral triangle from the vertices; to find the side of triangle. [Perkins' *Geometry*, Olney's *Geometry*.]

CALCULUS.

189. Proposed by J. E. SANDERS, Hackney, Ohio.

Solve $d^2y/dx^2 = -\beta^2(p+y)$, p and β being constants. The initial conditions are $y=0$ for $x=0$, l ; $dy/dx=0$ for $x=l/2$. [Merriman's *Mechanics*, 9th Ed., 1903, §62.]

190. Proposed by SAUL EPSTEIN, The University of Chicago, Chicago, Ill.

$$\int_0^\infty \frac{\sin x \cos \beta x}{x} dx, \int_0^\infty \frac{\sin x x \cos x}{x}.$$

MECHANICS.

170. Proposed by M. E. GRÄBER, A. M., Heidelberg University, Tiffin, Ohio.

Prove that the moment of inertia of an ogival head rotating about its geometrical axis is $\frac{\pi w}{g} \int_0^{R/(4n-1)} y^4 dx$, where w is the weight in pounds of a cubic foot of material, R the radius of the base of the ogive, and n the diameter of projectile.

DIOPHANTINE ANALYSIS.

123. Proposed by L. E. DICKSON, Ph. D., The University of Chicago.

Of two numbers $a_i b_i c_i d_i e_i$ ($i=1, 2$) it is given that their 10 digits a_1, \dots, e_2 form a permutation of 0, 1, ..., 9, and that the sum of the two is $x8951$. Give an immediate evaluation of x ; also list the possible pairs $a_1, a_2; \dots; e_1, e_2$.